

Available online at www.sciencedirect.com

J. Math. Anal. Appl. 340 (2008) 1380–1384

Journal of
**MATHEMATICAL
 ANALYSIS AND
 APPLICATIONS**

www.elsevier.com/locate/jmaa

New results of periodic solutions for a kind of Duffing type p -Laplacian equation

Yi Tang^{a,*}, Yaqiong Li^b^a *Department of Mathematics and Physics, Hunan First Normal College, Changsha 410002, PR China*^b *College of Mathematics and Econometrics, Hunan University, Changsha, Hunan 410082, PR China*

Received 21 April 2007

Available online 10 October 2007

Submitted by J. Mawhin

Abstract

By using Mawhin–Manásevich continuation theorem, some new sufficient conditions for the existence and uniqueness of periodic solutions of Duffing type p -Laplacian differential equation are established, which are complement of previously known results.

© 2007 Elsevier Inc. All rights reserved.

Keywords: p -Laplacian; Periodic solutions; Duffing equation; Topological degree

1. Introduction

In the last several years, the existence of periodic solutions for the Duffing equation, Rayleigh equation and Liénard type equation has been received a lot of attention. We refer the reader to [1–9] and the references cited therein. However, so far as we know, fewer papers discuss the existence and uniqueness of periodic solutions for Duffing type p -Laplacian differential equation. We only find that Zhang and Li [7] deal with the existence and uniqueness of periodic solutions for Duffing type p -Laplacian differential equation

$$(\varphi_p(x'(t)))' + Cx'(t) + g(t, x(t)) = e(t), \quad (1)$$

where $p > 1$ and $\varphi_p : R \rightarrow R$ is given by $\varphi_p(s) = |s|^{p-2}s$ for $s \neq 0$ and $\varphi_p(0) = 0$, C is a constant, g is a continuous function defined on R^2 and is T -periodic about t , e is a continuous periodic function defined on R with period T , $\int_0^T e(t) dt = 0$ and $T > 0$. Zhang and Li [7] provide a sufficient condition for such existence and uniqueness:

Theorem A 1. *Suppose that the following conditions (A_1) , (A_2) and (A_3) hold:*

(A_1) $(g(t, u_1) - g(t, u_2))(u_1 - u_2) < 0$ for $t, u_1, u_2 \in R$ and $u_1 \neq u_2$.

* Corresponding author.

E-mail address: yitang1972@yahoo.com.cn (Y. Tang).

(A₂) $xg(t, x) < 0$ for $|x| > 0$ and $t \in R$.

(A₃) There exist positive constants K and M such that

$$2^{2-p}MT^p < 1, \quad g(t, x) \geq -M|x|^{p-1} - K \quad \text{for } x \geq 0 \text{ and } t \in R.$$

Then Eq. (1) has a unique T -periodic solution.

However, upon examining their proof of Theorem A in [7], we have found that the condition (A₃) can be abandoned. In this paper, we will also discuss the existence and uniqueness of T -periodic solutions of Eq. (1). By using Mawhin–Manásevich continuation theorem, we establish some sufficient conditions for the existence and uniqueness of T -periodic solutions of Eq. (1). These results generalize and improve those in [7].

2. Main result

For convenience, define

$$C_T^1 := \{x \in C^1(R): x \text{ is } T\text{-periodic}\},$$

which is a Banach space endowed with the norm $\| \cdot \|$ defined by $\|x\| = |x|_\infty + |x'|_\infty$, for all x , and

$$|x|_\infty = \max_{t \in [0, T]} |x(t)|, \quad |x'|_\infty = \max_{t \in [0, T]} |x'(t)|.$$

For the T -periodic boundary value problem

$$(\varphi_p(x'(t)))' = f(t, x, x'), \quad (2)$$

where f is a continuous function and T -periodic in the first variable, we have the following result.

Lemma 1. (See [9].) Let B be the open in C_T^1 of center 0 and radius r . Assume that the following conditions hold:

- (i) For each $\lambda \in (0, 1)$ the problem $(\phi_p(x'))' = \lambda f(t, x, x')$ has no solution on the board of B .
- (ii) The continuous function F defined on R by $F(a) = 1/T \int_0^T f(t, a, 0) dt$ is such that $F(-r)F(r) < 0$.

Then (2) has at least one solution in \bar{B} .

By using Lemma 1, we obtain our main results:

Theorem 1. Let (A₁) hold. Suppose that there exists a constant $d \geq 0$ such that

(A₂^{*}) $xg(t, x) < 0$ for $|x| > d$ and $t \in R$.

Then Eq. (1) has a unique T -periodic solution.

Proof. Consider the homotopic equation of Eq. (1) as following:

$$(\varphi_p(x'(t)))' + \lambda Cx'(t) + \lambda g(t, x(t)) = \lambda e(t), \quad \lambda \in (0, 1). \quad (3)$$

By Lemma 2 in [7], together with (A₁), it is easy to see that Eq. (1) has at most one T -periodic solution. Thus, to prove Theorem 1, it suffices to show that Eq. (1) has at least one T -periodic solution. To do this, we shall apply Lemma 1. Firstly, we will claim that the set of all possible T -periodic solutions of Eq. (3) are bounded.

Let $x(t) \in C_T^1$ be an arbitrary solution of Eq. (3) with period T . By integrating two sides of Eq. (3) over $[0, T]$, and noticing that $x'(0) = x'(T)$, we have

$$\int_0^T g(t, x(t)) dt = 0. \quad (4)$$

As $x(0) = x(T)$, there exists $t_0 \in [0, T]$ such that $x'(t_0) = 0$, while $\varphi_p(0) = 0$ we see

$$|\varphi_p(x'(t))| = \left| \int_{t_0}^t (\varphi_p(x'(s)))' ds \right| \leq \lambda \int_0^T |C| |x'(t)| dt + \lambda \int_0^T |g(t, x(t))| dt + \lambda \int_0^T |e(t)| dt, \quad (5)$$

where $t \in [t_0, t_0 + T]$.

From (4), there exists $\bar{\xi} \in [0, T]$ such that

$$g(\bar{\xi}, x(\bar{\xi})) = 0.$$

In view of (A_2^*) , we obtain

$$|x(\bar{\xi})| \leq d.$$

Then, we have

$$|x(t)| = \left| x(\bar{\xi}) + \int_{\bar{\xi}}^t x'(s) ds \right| \leq d + \int_{\bar{\xi}}^t |x'(s)| ds, \quad t \in [\bar{\xi}, \bar{\xi} + T],$$

and

$$|x(t)| = |x(t - T)| = \left| x(\bar{\xi}) - \int_{t-T}^{\bar{\xi}} x'(s) ds \right| \leq d + \int_{t-T}^{\bar{\xi}} |x'(s)| ds, \quad t \in [\bar{\xi}, \bar{\xi} + T].$$

Combining the above two inequalities, we obtain

$$\begin{aligned} |x|_\infty &= \max_{t \in [0, T]} |x(t)| = \max_{t \in [\bar{\xi}, \bar{\xi} + T]} |x(t)| \\ &\leq \max_{t \in [\bar{\xi}, \bar{\xi} + T]} \left\{ d + \frac{1}{2} \left(\int_{\bar{\xi}}^t |x'(s)| ds + \int_{t-T}^{\bar{\xi}} |x'(s)| ds \right) \right\} \\ &\leq d + \frac{1}{2} \int_0^T |x'(s)| ds. \end{aligned} \quad (6)$$

Denote

$$E_1 = \{t: t \in [0, T], |x(t)| > d\}, \quad E_2 = \{t: t \in [0, T], |x(t)| \leq d\}.$$

Since $x(t)$ is T -periodic, multiplying $x(t)$ and (3) and then integrating it from 0 to T , in view of (A_2^*) , we get

$$\begin{aligned} \int_0^T |x'(t)|^p dt &= - \int_0^T (\varphi_p(x'(t)))' x(t) dt \\ &= \lambda \int_0^T g(t, x(t)) x(t) dt - \lambda \int_0^T e(t) x(t) dt \\ &= \lambda \int_{E_1} g(t, x(t)) x(t) dt + \lambda \int_{E_2} g(t, x(t)) x(t) dt - \lambda \int_0^T e(t) x(t) dt \\ &\leq \int_0^T \max\{|g(t, x(t))|: t \in R, |x(t)| \leq d\} |x(t)| dt + \int_0^T |e(t)| |x(t)| dt \\ &\leq DT |x|_\infty, \end{aligned} \quad (7)$$

where $D = \max\{|g(t, x(t))|: t \in R, |x(t)| \leq d\} + |e|_\infty$.

For $x(t) \in C(R, R)$ with $x(t+T) = x(t)$, and $0 < r \leq s$, by using Hölder inequality, we obtain

$$\left(\frac{1}{T} \int_0^T |x(t)|^r dt \right)^{1/r} \leq \left(\frac{1}{T} \left(\int_0^T (|x(t)|^r)^{\frac{s}{r}} dt \right)^{\frac{r}{s}} \left(\int_0^T 1 dt \right)^{\frac{s-r}{s}} \right)^{1/r} = \left(\frac{1}{T} \int_0^T |x(t)|^s dt \right)^{1/s},$$

this implies that

$$|x|_r \leq T^{\frac{s-r}{rs}} |x|_s \quad \text{for } 0 < r \leq s. \quad (8)$$

Then, in view of (6)–(8), we can get

$$\left(\int_0^T |x'(t)| dt \right)^p \leq T^{p-1} |x'(t)|_p^p = T^{p-1} \int_0^T |x'(t)|^p dt \leq T^{p-1} DT |x|_\infty \leq T^p D \left(d + \frac{1}{2} \int_0^T |x'(s)| ds \right). \quad (9)$$

Since $p > 1$, (9) yields that we can choose some positive constant M_1 such that

$$\int_0^T |x'(s)| ds \leq M_1, \quad |x|_\infty \leq d + \frac{1}{2} \int_0^T |x'(s)| ds \leq M_1.$$

In view of (5), we have

$$\begin{aligned} |x'|_\infty^{p-1} &= \max_{t \in [0, T]} \{ |\varphi_p(x'(t))| \} = \max_{t \in [t_0, t_0+T]} \left\{ \left| \int_{t_0}^t (\varphi_p(x'(s)))' ds \right| \right\} \\ &\leq \int_0^T |C| |x'(t)| dt + \int_0^T |g(t, x(t))| dt + \int_0^T |e(t)| dt \\ &\leq |C| M_1 + T \max \{ |g(t, x(t))| : t \in R, |x(t)| \leq M_1 \} + T |e|_\infty := M_1^*. \end{aligned} \quad (10)$$

Thus, we can get some positive constant $M_2 > M_1 + M_1^* + 1$ such that, for all $t \in R$,

$$|x'(t)| \leq M_2.$$

Hence taking $r = M_1 + M_2 + d + 1$, we have that the set of all possible T -periodic solutions of Eq. (3) is a subset of B . On the other hand, it is clear that, in our case $F(a) = -1/T \int_0^T g(t, a) dt$. From (A_2^*) it follows that $F(-r)F(r) < 0$. In consequence we can apply Manásevich–Mawhin continuation theorem to deduce that Eq. (3) has at least one solution in \tilde{B} . This completes the proof. \square

3. Examples and remarks

As an application, let us consider the following equation:

$$(\varphi_p x'(t))' + 100x'(t) - (67 + \cos^2 t)(x^9(t) - 12) = \cos t, \quad (11)$$

where $p = \sqrt{2}$. We can easily check the conditions (A_1) and (A_2^*) hold. By Theorem 1, Eq. (11) has a unique 2π -periodic solution.

Remark 1. Since $p = \sqrt{2}$, one can easily to see that neither [1–9] nor the references therein can obtain the existence and uniqueness of 2π -periodic solutions for (11). Abandoning the condition (A_3) , all the results in Ref. [7] are just the special cases with $d = 0$ of this paper. Moreover, we have corrected Eq. (7) in Ref. [7] as Eq. (5). To sum up, our research results are essentially new and the complement of previous known results.

Acknowledgments

This work was supported by the NNSF (10371034) of China. The authors are grateful to the referee for his or her suggestions on the first draft of the paper.

References

- [1] Xiankai Huang, Zigui Xiang, On the existence of 2π -periodic solution for delay Duffing equation $x''(t) + g(t, x(t - \tau)) = p(t)$, Chinese Sci. Bull. 39 (3) (1994) 201–203.
- [2] Yongkun Li, Periodic solutions of the Liénard equation with deviating arguments, J. Math. Res. Exposition 18 (4) (1998) 565–570 (in Chinese).
- [3] G.Q. Wang, S.S. Cheng, A priori bounds for periodic solutions of a delay Rayleigh equation, Appl. Math. Lett. 12 (1999) 41–44.
- [4] Genqiang Wang, Jurang Yan, On existence of periodic solutions of the Rayleigh equation of retarded type, Int. J. Math. Math. Sci. 23 (1) (2000) 65–68.
- [5] Shiping Lu, Weigao Ge, Some new results on the existence of periodic solutions to a kind of Rayleigh equation with a deviating argument, Nonlinear Anal. 56 (2004) 501–514.
- [6] Minggang Zong, Hongzhen Liang, Periodic solutions for Rayleigh type p -Laplacian equation with deviating arguments, Appl. Math. Lett. 206 (2007) 43–47.
- [7] Fuxing Zhang, Ya Li, Existence and uniqueness of periodic solutions for a kind of Duffing type p -Laplacian equation, Nonlinear Anal. Real World Appl. (2007), doi:10.1016/j.nonrwa.2007.01.013, in press.
- [8] R.E. Gaines, J. Mawhin, Coincidence Degree, and Nonlinear Differential Equations, Lecture Notes in Math., vol. 568, Springer-Verlag, Berlin, 1977.
- [9] R. Manásevich, J. Mawhin, Periodic solutions for nonlinear systems with p -Laplacian-like operators, J. Differential Equations 145 (1998) 367–393.